syngenta

FIFRA

# Section 24(c) Special Local Need Label

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF SOUTH DAKOTA**

**Flexstar®**

**For Control of Weeds in Soybeans in South Dakota West of Highway 281 in the Counties of Aurora, Beadle, Brown, Brule, Buffalo, Campbell, Charles Mix, Douglas, Edmunds, Faulk, Gregory, Hand, Hughes, Hyde, Jerauld, Lyman, McPherson, Potter, Spink, Sully, Tripp, Walworth**

**EPA Reg. No. 100-1101**
**EPA SLN No. SD-190004**

**This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2024**

**DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.
- Failure to follow the directions for use and precautions on this label may result in poor pest control, crop injury, or illegal residues.

**Application Methods**
Apply Flexstar at 0.75 pt/A as a postemergence broadcast application to soybeans. Apply in a minimum of 15 GPA.

**Spray Additives**
Add 1% v/v methylated seed oil (MSO) plus ammonium sulfate (AMS) at a minimum of 8.5 lbs/100 gals of finished spray volume.

**Weeds Controlled**

| Weed | Maximum Number of Leaves At Application |
|---|---|
| Cocklebur, common | 2 |
| Mustard, wild | 4 |
| Nightshade, black | 2 |
| Pigweed, redroot | 4 |
| Waterhemp, common | 2 |
| Smooth Pigweed | 2 |
| Common Ragweed | 4, (suppression only) |
| Palmer Amaranth, | 2 |
| Common Lambsquarter | 2 (suppression only) |

**Rotational Crop Restrictions**

| Rotational Crops | Minimum Rotational Interval after Flexstar Application (Months) |
|---|---|
| Bean, Dry<br>Potato<br>Soybean | 0 months |
| Small Grains, such as Wheat, Barley, Rye | 9 months |
| Edible-podded beans and peas not otherwise specified in this label<br>Pea, Dry | 12 months |
| Corn, Field*<br>Corn, Seed*<br>Sunflowers*<br>Grain and Forage Sorghum* | 18 months |
| All other crops not listed above | Minimum 18 months and a successful field bioassay (refer to **Field Bioassay Instructions** section) must be performed before planting rotational crop |

*For fields with greater than 4% organic matter, a successful field bioassay (refer to **Field Bioassay Instructions** section) must be performed before planting field corn or seed corn.

**Field Bioassay Instructions**

Plant several strips of the desired crop variety across the field which has been previously treated with Flexstar. Plant the strips perpendicular to the direction Flexstar was applied. The strips should be located so that all the different field conditions are encountered, including differences in soil texture, organic matter, pH, and drainage. If the crop does not show visible symptoms of injury and/or stand reduction, this field can be planted to this crop. If visible injury and/or stand reduction occur, do not plant the field to this crop; plant only a labeled crop.

**Use Restrictions**

- Do not exceed 0.75 pt/A (0.176 lb ai/A) of Flexstar per acre in any one year. Do not apply to any field more than once every two years.
- Do not make a Flexstar application after June 20.
- Do not harvest soybeans within 45 days of application.
- Avoid overlapping spray swaths, as injury may occur to rotational crops.

---

©2019 Syngenta

Flexstar® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: SD1101019AA0319